

MAT 167 TEST 1 FORM B (LIMITS)

Directions: Solve each required problem **on a separate sheet of paper**. Use pencil and show all work; I deduct points for using pen or skipping important steps. **You must shut off your cell phone**. Some problems are worth more than others. Take your time; **quality is preferred to quantity**. I encourage you to ask questions.


1. Give (a) an intuitive definition of continuity of a function f at a point $x = a$, and (b) the precise definition of continuity of a function f at a point $x = a$.

Solution: f is continuous at a point $x = a$ if

- (a) its graph has no holes, skips, or vertical asymptotes.
 (b) we can find the limit by substitution; that is, $\lim_{x \rightarrow a} f(x) = f(a)$.
-

2. Graph the function $f(x) = x^2 - 4x + 3$. Identify the point $(a, f(a))$ at which the function has a tangent line with zero slope, and confirm your answer by making a table of slopes of secant lines to approximate the slope of the tangent line at this point.

Solution: The graph of $f(x) = x^2 - 4x + 3$ is at right. It looks as if the function has a tangent line with zero slope at $x = 3$.



x	1	1.9	1.99	1.999	2.001	2.01	2.1	3
$\frac{f(x)-f(2)}{x-2}$	-1	-0.1	-0.01	-0.001	0.001	0.01	0.1	1

It looks as if the slope of the tangent line will be 0.

3. Use a table of values to estimate $\lim_{x \rightarrow 0^+} (x \ln x)$, correct to two decimal places.

Solution: I would use the table below, but there are other possibilities.

x	1	0.1	0.01	0.001	10^{-4}	10^{-5}	10^{-6}
$x \ln x$	0	-0.2303	-0.0461	-0.0069	-0.0009	10^{-4}	10^{-5}

(You don't have to use the 10^{-n} notation. I did so in order to save time.)

4. If possible, determine a value of b such that $\lim_{x \rightarrow 2} p(x)$ exists.

$$p(x) = \begin{cases} 3x + b & \text{if } x \leq 2 \\ x - 2 & \text{if } x > 2. \end{cases}$$

Solution: We need $\lim_{x \rightarrow 2^-} p(x) = \lim_{x \rightarrow 2^+} p(x)$. For the left-sided limit, $x < 2$, so we use $p(x) = 3x + b$. For the right-sided limit, $x > 2$, so we use $p(x) = x - 2$. By substitution,

$$\begin{aligned}\lim_{x \rightarrow 2^-} (3x + b) &= \lim_{x \rightarrow 2^+} (x - 2) \\ 6 + b &= 0 \\ b &= -6.\end{aligned}$$

5. The table below gives various values of three functions f, g, h around the point $x = \pi$.
- What would be the value of $\lim_{x \rightarrow \pi} f(x)$ if we applied the Squeeze Theorem?
 - What aspects of the data suggest that we can apply the Squeeze Theorem?
 - What additional information would we need to be sure that we can?

x	2.5	3	3.1	$\lim_{x \rightarrow \pi} \dots$	3.2	3.3	3.5
$g(x)$	11	7	6	5.8	5.5	5.3	5.1
$f(x)$	10	-5	6	?	5	-1	4
$h(x)$	-20	-10	0	5.8	5	-3	-12

Solution: (a) 5.8

(b) It *looks* as if $h(x) \leq f(x) \leq g(x)$ for all x in the domain $[2.5, 3.5]$.

(c) Since the table only gives a sample of six points out of infinitely many, we would need to know that f really does lie between g and h for all $x \in [2.5, 3.5]$ — or at least for some neighborhood of π .

6. Determine the interval(s) on which the function $g(x) = \sqrt{x^4 - 1}$ is continuous.

Solution: We know from class that this type of function — the composition of two functions that are continuous on their domains — is continuous on its domain. So, we need merely find the domain. A number inside a square root should be nonnegative:

$$\begin{aligned}x^4 - 1 &\geq 0 \\ x^4 &\geq 1.\end{aligned}$$

This is true if $x \geq 1$ or if $x \leq -1$. Hence, g is continuous on $(-\infty, -1] \cup [1, \infty)$.

7. Identify and classify (as holes, jumps, or vertical asymptotes) all the discontinuities of

$$r(x) = \frac{x^4 + 7}{x^5 + x^2 - x}.$$

Also find the horizontal asymptotes. Do not use any shortcuts you may have learned in pre-calculus, unless you're willing to explain them using principles of Calculus.

Solution: We know from class that this function — a *rational* function — is continuous on its domain. It is defined everywhere except where the denominator is zero. That is,

$$\begin{aligned}x^5 + x^2 - x &= 0 \\x(x^4 + x - 1) &= 0 \\x &= 0, -1.22074, 0.72449.\end{aligned}$$

Hence, it is discontinuous at each of these points. At each of these three values, the numerator is nonzero; in fact, the numerator is always positive. Hence, all three are vertical asymptotes.

We find the horizontal asymptote by considering

$$\lim_{x \rightarrow \infty} r(x) = \lim_{x \rightarrow \infty} \frac{x^4 + 7}{x^5 + x^2 - x} = \lim_{x \rightarrow \infty} \left(\frac{x^4 + 7}{x^5 + x^2 - x} \cdot \frac{\frac{1}{x^4}}{\frac{1}{x^4}} \right) = \lim_{x \rightarrow \infty} \frac{1 + \frac{7}{x}}{x + \frac{1}{x^2} - \frac{1}{x^3}} = 0.$$

Note: Apparently, I made a typographical error when writing this problem, which requires you to use an approximation technique to compute all the roots. You shouldn't encounter a similar problem with the function I give you on the exam.

8. List the first five terms of the sequence $\frac{n^2 + 1}{n}$, then find $\lim_{n \rightarrow \infty} \frac{n^2 + 1}{n}$.

Solution: The first five terms are

$$\frac{1+1}{1} = 2, \quad \frac{4+1}{2} = \frac{5}{2}, \quad \frac{9+1}{3} = \frac{10}{3}, \quad \frac{16+1}{4} = \frac{17}{4}, \quad \frac{25+1}{5} = \frac{26}{5}.$$

The limit is

$$\lim_{n \rightarrow \infty} \left(\frac{n^2 + 1}{n} \cdot \frac{\frac{1}{n}}{\frac{1}{n}} \right) = \lim_{n \rightarrow \infty} \frac{n + \frac{1}{n}}{1} = \infty.$$

9. If $f(x) = 4x - 3$, find

- $L = \lim_{x \rightarrow 3} f(x)$ using the properties of limits; and
- δ such that if $|x - 3| < \delta$, then $|f(x) - L| < 0.01$.

Solution: (a) Since f is a polynomial, we can substitute: $f(x) = 4 \cdot 3 - 3 = 9$. We can also use more basic properties:

$$\begin{aligned}\lim_{x \rightarrow 3} (4x - 3) &= \lim_{x \rightarrow 3} (4x) - \lim_{x \rightarrow 3} 3 \text{ (sum property of limits)} \\&= 4 \lim_{x \rightarrow 3} x - \lim_{x \rightarrow 3} 3 \text{ (constant multiple property)} \\&= 4 \cdot 3 - \lim_{x \rightarrow 3} 3 \text{ (limit of powers of } x\text{)} \\&= 12 - 3 \text{ (limit of a constant)} \\&= 9.\end{aligned}$$

(b) We need

$$|f(x) - 9| < 0.01$$

$$|(4x - 3) - 9| < 0.01$$

$$|4x - 12| < 0.01$$

$$4|x - 3| < 0.01$$

$$|x - 3| < 0.0025 = \frac{1}{400}.$$

The left side has the form $|x - 3| < \delta$! Thus, if we set $\delta = \frac{1}{400}$, then $|f(x) - L| < 0.01$.
